

Pr imbruvica[®] EXPERIENCE

 **AVAILABLE IN
CANADA SINCE 2014**

12 CLINICAL TRIALS
conducted across indications¹

>177,000 PERSON-YEARS
of cumulative global exposure,
combined across all indications^{1*}

9
INDICATIONS²

CLL				MCL	WM		MZL	cGVHD
Previously treated		Previously untreated active			Monotherapy	I [†] + R [§]		
Monotherapy	I [†] + BR [‡]	Monotherapy	I [†] + G [¶]					
2014	2017	2016	2019	2017	2016	2019	2018	2017

† I=IMBRUVICA[®]; ‡ BR=bendamustine-rituximab; § R=rituximab; ¶ G=obinutuzumab.

IMBRUVICA[®] (ibrutinib) is indicated:

- For the treatment of patients with previously untreated active chronic lymphocytic leukemia (CLL), including those with 17p deletion.
Clinical effectiveness of IMBRUVICA[®] in previously untreated patients with CLL with 17p deletion is based on the benefit observed in patients with CLL with 17p deletion who have received at least one prior therapy. Clinical trial data in previously untreated patients with CLL with 17p deletion are very limited.
- In combination with obinutuzumab for the treatment of patients with previously untreated active CLL, including those with 17p deletion.
- For the treatment of patients with CLL who have received at least one prior therapy, including those with 17p deletion.
- In combination with bendamustine and rituximab for the treatment of patients with CLL who have received at least one prior therapy.

Clinical trial data with IMBRUVICA[®] in combination with bendamustine and rituximab in patients with CLL with 17p deletion are limited.

- For the treatment of patients with relapsed or refractory mantle cell lymphoma (MCL).
- For the treatment of patients with marginal zone lymphoma (MZL) who require systemic therapy and have received at least one prior anti-CD20-based therapy.
- For the treatment of patients with Waldenström's macroglobulinemia (WM).
Clinical effectiveness of IMBRUVICA[®] is based on response rates demonstrated in a single-arm study in patients who had received at least one prior therapy.
- In combination with rituximab for the treatment of patients with WM.
- For the treatment of patients with steroid dependent or refractory chronic graft versus host disease (cGVHD).

Consult the Product Monograph at <http://www.janssen.com/canada/products> for information regarding warnings, precautions, adverse reactions, interactions, dosing and conditions of clinical use. The Product Monograph is also available by calling 1-800-567-3331 or 1-800-387-8781.

* From January 2014 (launch) to October 31, 2019.

References:

1. Data on file, Janssen Inc. 2. IMBRUVICA[®] Product Monograph, Janssen Inc., April 17, 2020.

All trademarks used under license. I IMBRUVICA[®] is co-developed with Pharmacyclics. Janssen Inc. is the marketing authorization holder and is the responsible editor of this document. © 2020 Pharmacyclics I © 2020 Janssen Inc., 19 Green Belt Drive, Toronto, ON M3C 1L9 www.janssen.com/canada | CP-162724E

MEMBER OF
INNOVATIVE MEDICINES CANADA



pharmacyclics[®]
An AbbVie Company

janssen
PHARMACEUTICAL COMPANIES OF
johnson & johnson

imbruvica[®]
(ibrutinib) capsules